




**Figure 1. Troxatyl + STI - KBM-5**




**Figure 2. Troxatyl + STI - KBM-5R**




**Figure 3. Troxatyl + STI-KBM-7**




**Figure 4. Troxatyl + STI-KBM-7R**




**Figure 5. Troxatyl + STI-KBM-5R; Caspase 3/7 activity-48 h (2 h incubation)**




**Figure 6. Troxatyl + STI-KBM-5R; Caspase 3/7 activity-72 h (2 h incubation)**



**Figure 7. Troxatyl + STI-KBM-7R; Caspase 3/7 activity  
h (2 h incubation)**




**Figure 8. Troxatyl + STI-KBM-7R; Caspase 3/7 activity 10 h (2 h incubation)**






**Figure 9. Troxatyl + STI-KBM-7R; Caspase 3/7 activity 27 h (2 h incubation)**



**Figure 10. Troxatyl + STI-KBM-7R; Caspase  
3/7 activity-48h (2 h incubation)**



**Figure 11. Troxatyl + STI-KBM-7R; Caspase  
3/7 activity-72h (2 h incubation)**




Figure 12. Survival of SCID mice bearing KBM-5 cells




Figure 13. Survival of SCID mice bearing KBM-5R cells




Figure 14. Survival of SCID mice bearing KBM-5 cells

